

Leonard M. Parver, M.D.
Robert F. Stephens, M.D.
David L. Parver, M.D.
Jordan L. Heffez, M.D.
Fadi P. Nasrallah, M.D.
Nishi Gulati, M.D.

Washington, DC 1145 Nineteenth St., NW Suite 500 Washington, DC 20036 T: 202.833.1668 F: 202.833.4698 **Bethesda, MD**6410 Rockledge Dr.
Suite 400
Bethesda, MD 20817
T: 301.530.5200
F: 301.493.6577

Annandale, VA 3289 Woodburn Rd. Suite 270 Annandale, VA 22003 T: 703.849.8601 F: 703.849.8605

INFORMATION REGARDING DILATING EYE DROPS

Dilating drops are used to dilate or enlarge the pupils of the eye to allow the ophthalmologist to get a better view of the inside of your eye.

Dilating drops frequently blur vision for a length of time which varies from person to person and may make bright lights bothersome. It is not possible for your ophthalmologist to predict how much your vision will be affected. Because driving may be difficult immediately after an examination, it's best if you make arrangements not to drive yourself.

Adverse reaction, such as acute angle-closure glaucoma, may be triggered from the dilating drops. This is extremely rare and treatable with immediate medical attention.

I hereby authorize Drs. Stephens, L. Parver, Nasrallah, D. Parver, J. Heffez, N. Gulati and/or such assistants as may be designated by him/her to administer dilating eye drops. The eye drops are necessary to diagnose my condition.

| Patient (or person authorized to sign for patient) | Date | |
|--|------|--|
| | | |
| Witness | Date | |